





BEST AVAILABLE COPY



Medline

[Simple](#) | [Advanced](#) | [Citation](#) | [History](#) | [Results](#) | [Record](#)

Alterations in the biosynthesis of lignin in transgenic plants with chimeric genes for 4-coumarate: coenzyme A ligase.

Kajita S, Katayama Y, Omori S
Plant Cell Physiol 1996 Oct 37:957-65

BROWSE : [Plant Cell Physiol](#) • [Volume 37](#) • [Issue 7](#)

VIEW : [MEDLINE](#), [full MEDLINE](#), [related records](#)

Abstract

The introduction of chimeric sense and antisense gene constructs for 4-coumarate:coenzyme A ligase into tobacco plants caused the reduction of the 4CL activity in the transgenic plants. In the transgenic plants, the cell walls of the xylem tissue in stems were brown and the molecular structure of lignin in the colored cell walls was dramatically different from that in the control plants. Analysis with different types of stain revealed that levels of cinnamyl aldehyde residues and syringyl units in lignin were depressed in the brownish cell walls. Furthermore, the lignin content in colored tissue was lower than that in the normal tissue. Our results indicate that 4CL has important roles in the determination of the composition and the amount of lignin in tobacco plants.

MeSH

[Blotting, Northern](#); [Chimeric Proteins](#); [Coenzyme A Ligases](#); [Lignin](#); [Molecular Sequence Data](#); [Plants, Transgenic](#); [Tobacco](#)

Author Address

Research Center, Mitsubishi Paper Mills Ltd., Ibaraki, Japan.

[Order Document](#)

BioMedNet

Research
Tools

Reviews

Journal
CollectionNews &
CommentBooks &
LabwareScience
Jobs
[Information for Advertisers](#)

© Elsevier Science Limited 2000